

# Hedging in complete markets driven by normal martingales

Youssef El-Khatib and Nicolas Privault

*Département de Mathématiques, Université de La Rochelle  
Avenue Michel Crépeau, 17042 La Rochelle Cedex 1, France\**

February 4, 2003

## Abstract

This paper aims to a unified treatment of hedging in market models driven by martingales with deterministic bracket  $\langle M, M \rangle_t$ , including Brownian motion and the Poisson process as particular cases. Replicating hedging strategies for European, Asian and Look-back options are explicitly computed using either the Clark-Ocone formula or an extension of the delta hedging method, depending on which is most appropriate.

**Key words:** Normal martingales, chaos representation property, hedging strategies, exotic options.

*Classification:* 90A09, 90A12, 90A60, 60H07.

## Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b>Notation and preliminaries</b>	<b>4</b>
2.1	Chaotic calculus . . . . .	4
2.2	Structure equations . . . . .	5
2.3	Change of variable formula . . . . .	8
2.4	Girsanov theorem . . . . .	9
<b>3</b>	<b>Computations of predictable representations</b>	<b>10</b>
3.1	Clark formula . . . . .	10
3.2	Deterministic structure equation . . . . .	10
3.3	Markovian case . . . . .	13

---

\*yelkhati@univ-lr.fr, nprivaul@univ-lr.fr

<b>4</b>	<b>Computations of hedging strategies</b>	<b>14</b>
4.1	Market model . . . . .	14
4.2	European options and deterministic structure . . . . .	16
4.3	Asian options and deterministic structure . . . . .	19
4.4	European options and Azéma martingales . . . . .	22
4.5	Lookback options . . . . .	23

# 1 Introduction

The Clark formula [Cla70] allows in principle for the calculation of replicating hedging strategies in complete markets [KO91], however explicit computations are in general difficult to perform via this formula. For markets driven by Brownian motion a proof of the classical Black-Scholes formula via the Clark-Ocone formula can be found in [Øks96], Ch. 5, p. 13. This method has been recently extended to markets driven by a Poisson process in [AOPU00]. Brownian motion and the compensated Poisson process share the important chaos representation property which is capital for market completeness.

In this paper we consider a larger family of martingales  $(M_t)_{t \in [0, T]}$  satisfying the following two conditions:

- a) the chaotic representation property (in relation with market completeness), i.e. every square-integrable functional, measurable with respect to the filtration generated by  $(M_t)_{t \in [0, T]}$ , can be expanded into a series of multiple stochastic integrals of deterministic functions with respect to  $(M_t)_{t \in [0, T]}$ ,
- b) the condition  $d\langle M, M \rangle_t = \alpha_t^2 dt$ , where  $(\alpha_t)_{t \in [0, T]}$  is a square integrable deterministic function.

Hypothesis (b) implies that  $([M, M]_t - \int_0^t \alpha_s^2 ds)_{t \in [0, T]}$  is a martingale, hence from (a) there exists a process  $(\phi_t)_{t \in [0, T]}$  such that  $(M_t)_{t \in [0, T]}$  satisfies the structure equation

$$d[M, M]_t = \alpha_t^2 dt + \phi_t dM_t, \quad t \in [0, T], \tag{1.0.1}$$

cf. [É90]. This equation can be viewed as a decomposition of  $d[M, M]_t$  by projection on  $dt$  and  $dM_t$ , which yields a closed Itô type change of variable formula, see Prop. 2.3.

Brownian motion is obtained for  $\alpha_t = 1$  and  $\phi_t = 0$ ,  $t \in [0, T]$ , and the Poisson process corresponds to non zero constant  $\phi_t$ ,  $t \in [0, T]$ . The choice

$\phi_t = \beta M_t$ ,  $-2 \leq \beta < 0$ , considered in [DP99], corresponds to the Azéma martingale and yields another complete market model with jumps. Choosing  $(\phi_t)_{t \in [0, T]}$  to be a deterministic function allows for the driving process to be alternatively Brownian or Poisson, depending on the vanishing of  $\phi_t$ , see [JP02] for the corresponding market model.

The Clark-Ocone formula states the predictable representation of a random variable  $F$  as

$$F = E[F] + \int_0^T E[D_t F \mid \mathcal{F}_t] dM_t,$$

where  $(\mathcal{F}_t)_{t \in [0, T]}$  is the filtration generated by  $(M_t)_{t \in [0, T]}$  and  $D_t$  is the gradient operator that lowers the degree of multiple stochastic integrals with respect to  $(M_t)_{t \in [0, T]}$ . One of the goals of this paper is to compute the process  $t \mapsto E[D_t F \mid \mathcal{F}_t]$  in several situations. We obtain explicit hedging formulas for European calls in the mixed Brownian-Poisson model of [JP02] and in the Azéma martingale model of [DP99] and for Asian and Lookback options.

More precisely, let  $(S_{t, T}^x)_{t \in [0, T]}$  denote the stock price process driven by  $(M_t)_{t \in [0, T]}$ , starting from  $x$  at time  $t$ , with volatility  $(\sigma_t)_{t \in [0, T]}$ , and let  $i_t = 1_{\{\phi_t=0\}}$ ,  $j_t = 1 - i_t$ ,  $S_t = S_{0, t}^1$ ,  $t \in [0, T]$ . In a model with deterministic structure equation, i.e.  $d[M, M]_t = \alpha_t^2 dt + \phi_t dM_t$  with deterministic  $(\phi_t)_{t \in [0, T]}$ , the replicating hedging strategy of a European call with payoff  $(S_T - K)^+$  is given by

$$t \mapsto \frac{e^{-\int_t^T r_s ds}}{\sigma_t S_t} E \left[ i_t \sigma_t S_{t, T}^x 1_{\{S_{t, T}^x \geq K\}} + \frac{j_t}{\phi_t} (\sigma_t \phi_t S_{t, T}^x - (K - S_{t, T}^x)^+) 1_{\{S_{t, T}^x \geq \frac{K}{1 + \sigma_t}\}} \right]_{x=S_t},$$

cf. (4.1.6) and Prop. 4.1. This formula extends both the classical Black-Scholes hedging formula in the Brownian case ( $\phi_t = 0$ ,  $t \in [0, T]$ ), and the hedging formula of [AOPU00], Th. 6.1, in the Poissonian jump case ( $\phi_t = 1$ ,  $t \in [0, T]$ ). It can be obtained both from the Clark formula and from martingale methods. The above conditional expectation can be also explicitly computed, see. Prop. 4.2. The case of Asian options is treated in Prop. 4.3 in the deterministic structure equation model, and Lookback options are considered in Prop. 4.5 in a market driven by Brownian motion. In the Azéma martingale model of [DP99], i.e.  $d[M, M]_t = dt + \beta M_t^- dM_t$ ,

$-2 \leq \beta < 0$ , we obtain

$$t \mapsto \frac{e^{-\int_t^T r_s ds}}{\beta M_t \sigma_t S_t} E \left[ \sigma_t \beta (y + M_T - M_t) S_{t,T}^x 1_{[\frac{K}{1+\sigma_t \beta (y+M_T-M_t)}, \infty)}(S_{t,T}^x) \right. \\ \left. + (S_{t,T}^x - K) 1_{[\frac{K}{1+\sigma_t \beta (y+M_T-M_t)}, K]}(S_{t,T}^x) \right]_{x=S_t}^{y=M_t},$$

see Prop. 4.4.

We proceed as follows. In Sect. 2 we introduce the notation of chaotic calculus, the solutions of structure equations and the Clark-Ocone formula which gives the predictable representation of the random variable  $F$ . We also state the change of variable formula and Girsanov theorem which hold in a particular form for solutions of structure equations. In Sect. 3 we describe different methods for the computation of predictable representations for the general class of normal martingales having the chaos representation property. The intrinsic expression of the gradient  $D$  is completely known if  $(\phi_t)_{t \in [0, T]}$  is deterministic, i.e. for the Brownian, Poisson and deterministic structure equation models, (Sect. 3.2). In the Markovian case (Sect. 3.3) it is possible to combine the Clark-Ocone and Itô formulas to obtain the explicit predictable representation of  $F$ . Sect. 4 is devoted to the computation of replicating portfolios. In Sect. 4.2 we hedge European calls using the Clark formula, extending the method applied in the Brownian case in [Øks96], Ch. 5, pp. 13-15. In Sects. 4.3, 4.4 and 4.5 we deal with Asian, European and Lookback options, in particular we use the delta hedging approach to recover some results obtained in [Ber98] from the Clark formula.

## 2 Notation and preliminaries

### 2.1 Chaotic calculus

Let  $(M_t)_{t \in [0, T]}$  be a martingale on the space  $\Omega$  of càdlàg functions from  $[0, T]$  to  $\mathbb{R}$ , having the chaos representation property. Let  $(\alpha_t)_{t \in [0, T]} \in L^2([0, T])$  be a positive deterministic non vanishing square-integrable function, and assume that  $(M_t)_{t \in [0, T]}$  has deterministic angle bracket  $d\langle M, M \rangle_t = \alpha_t^2 dt$ . We denote by  $(\mathcal{F}_t)_{t \in [0, T]}$  the filtration generated by  $(M_t)_{t \in [0, T]}$ , and by  $L_\alpha^2([0, T])^{on}$  the space  $L^2([0, T], \alpha_t^2 dt)^{on}$  of  $\alpha_{t_1}^2 \cdots \alpha_{t_n}^2 dt_1 \cdots dt_n$  square-integrable symmetric functions. The multiple stochastic integral  $I_n(f_n)$  is defined as

$$I_n(f_n) = n! \int_0^T \int_0^{t_n} \cdots \int_0^{t_2} f_n(t_1, \dots, t_n) dM_{t_1} \cdots dM_{t_n}, \quad n \geq 1,$$

$f_n \in L^2_\alpha([0, T])^{\circ n}$ , with

$$E[I_n(f_n)I_m(g_m)] = n!1_{\{n=m\}}\langle f_n, g_m \rangle_{L^2_\alpha([0, T])^{\circ n}}. \quad (2.1.1)$$

The chaos representation property for  $(M_t)_{t \in [0, T]}$  states that every  $F \in L^2(\Omega)$  has a decomposition as  $F = \sum_{n=0}^\infty I_n(f_n)$ . Let  $D : \text{Dom}(D) \rightarrow L^2(\Omega \times [0, T], dP \times \alpha_t^2 dt)$  denote the closable, unbounded gradient operator defined as

$$D_t F = \sum_{n=1}^\infty n I_{n-1}(f_n(*, t)), \quad dP \times dt - a.e.,$$

with  $F = \sum_{n=0}^\infty I_n(f_n)$  in  $\text{Dom}(D)$ , i.e.

$$\sum_{n=1}^\infty nn! \|f_n\|_{L^2(\mathbb{R}_+^n)}^2 < \infty.$$

## 2.2 Structure equations

Let  $L_{ad}^\infty(\Omega \times [0, T])$  be the space of bounded,  $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$ -adapted stochastic processes. We assume that  $(M_t)_{t \in [0, T]}$  is solution of the structure equation

$$d[M, M]_t = \alpha_t^2 dt + \phi_t dM_t, \quad t \in [0, T], \quad (2.2.1)$$

where  $\phi_t = \varphi(t, M_{t-})$  is a deterministic function of  $t$  and  $M_t$ . Existence and uniqueness of the solutions are guaranteed when  $\phi_t$  is a deterministic function [É90]. Existence is proved when  $\phi_t = \varphi(M_{t-})$  and  $\varphi$  is a continuous function [Mey89], and the solution is unique when  $\varphi(x) = \beta x$  with  $\beta \in [-2, 0)$ , cf. [É90]. See also [Pha00], [Tav99] for recent results on structure equations. Let  $i_t = 1_{\{\phi_t=0\}}$  and  $j_t = 1_{\{\phi_t \neq 0\}} = 1 - i_t$ ,  $t \in [0, T]$ . The continuous part of  $(M_t)_{t \in [0, T]}$  is given by  $dM_t^c = i_t dM_t$  and the eventual jump of  $(M_t)_{t \in [0, T]}$  at time  $t \in [0, T]$  is given as  $\Delta M_t = \phi_t$ ,  $t \in [0, T]$ , see [É90], p. 77.

- a) If  $(\phi_t)_{t \in [0, T]}$  is deterministic, let  $\lambda_t = j_t \alpha_t^2 / \phi_t^2$ ,  $t \in [0, T]$ , then  $(M_t)_{t \in [0, T]}$  can be represented as

$$dM_t = i_t \alpha_t dB_t + \phi_t (dN_t - \lambda_t dt), \quad t \in [0, T], \quad M_0 = 0, \quad (2.2.2)$$

where  $(B_t)_{t \in [0, T]}$  is a standard Brownian motion, and  $(N_t)_{t \in [0, T]}$  a Poisson process independent of  $(B_t)_{t \in [0, T]}$ , with intensity  $\nu_t = \int_0^t \lambda_s ds$ ,  $t \in [0, T]$ , cf. Prop. 4 of [É90], and [JP02].

b) If  $\phi_t = \beta M_t$ ,  $\beta \in \mathbb{R}$ , then (2.2.1) has a unique solution called the Azéma martingale, cf. [É90]. If  $-2 \leq \beta < 0$ , this solution has the chaos representation property and it has been used to model a complete market with jumps in [DP99].

Figure 1 shows a sample path of  $(S_t)_{t \in [0, T]}$  and the corresponding function  $(i_t)_{t \in [0, T]}$  chosen to be a simple indicator function, with  $S_0 = 4$ ,  $\sigma_t = 1$ ,  $\alpha_t^2 = 50$ ,  $\phi_t = 1.6i_t$ ,  $t \in [0, T]$ .




Figure 1 - Sample trajectory of  $(S_t)_{t \in [0, T]}$ \*

Figure 2 is a simulation of an Azéma martingale with  $\beta \neq -1$ , from a discretization of the structure equation (2.2.1):

$$\Delta X_t = \frac{\beta X_{t-}}{2} \pm \sqrt{\left(\frac{\beta X_{t-}}{2}\right)^2 + \Delta t},$$

with probabilities

$$p = \frac{1}{2} \mp \frac{\beta X_{t-}}{\sqrt{\left(\frac{\beta X_{t-}}{2}\right)^2 + \Delta t}}.$$

---

\*Vertical lines represent jumps.




Figure 2 - Sample path of an Azéma martingale\*

In all cases of interest in this paper we have  $(\phi_t)_{t \in [0, T]} \in L_{ad}^\infty(\Omega \times [0, T])$ , since if  $(\phi_t)_{t \in [0, T]} = (\beta M_t)_{t \in [0, T]}$  then  $\sup_{t \in [0, T]} |M_t| \leq (-2/\beta)^{1/2}$  (see [É90], p. 83). Given  $(u_t)_{t \in [0, T]} \in L_{ad}^\infty(\Omega \times [0, T])$ , we denote by  $(\xi_t(u))_{t \in [0, T]}$  the solution of the equation

$$Z_t = 1 + \int_0^t Z_{s-} u_s dM_s, \quad t \in [0, T], \quad (2.2.3)$$

which can be written as ([Pro90], Th. 36, p. 77):

$$\xi_t(u) = \exp \left( \int_0^t u_s dM_s - \frac{1}{2} \int_0^t u_s^2 \alpha_s^2 ds \right) \prod_{s \in J_M^t} (1 + u_s \phi_s) e^{-u_s \phi_s}, \quad (2.2.4)$$

where  $J_M^t$  denotes the set of jump times of  $(M_s)_{s \in [0, t]}$ ,  $t \in [0, T]$ , and let  $\xi(u) = \xi_T(u)$ . If  $u \in L^\infty([0, T])$  then  $\xi_t(u)$  can be represented as

$$\xi_t(u) = \sum_{n=0}^{\infty} \frac{1}{n!} I_n((u1_{[0, t]})^{\otimes n}),$$

and we have  $D_s \xi_t(u) = 1_{[0, t]}(s) u_s \xi_t(u)$ ,  $s, t \in [0, T]$ .

**Definition 2.1** *Let  $\mathcal{S}$  denote the linear space generated by exponential vectors of the form  $\xi(u)$ , where  $u \in L^\infty([0, T])$ .*

The space  $\mathcal{S}$  is dense in  $L^2(\Omega)$ , and from the following Lemma  $\mathcal{S}$  is an algebra for the pointwise multiplication of random variables if  $(\phi_t)_{t \in [0, T]}$  is deterministic. The following is a version of Yor's formula [Yor76] or Th. 37 of [Pro90], page 79, for martingales with deterministic bracket  $\langle M, M \rangle_t$ .

**Lemma 2.2** For any  $u, v \in L^\infty([0, T])$ , we have the relation

$$\xi(u)\xi(v) = \exp(\langle u, v \rangle_{L_\alpha^2([0, T])})\xi(u + v + \phi uv). \quad (2.2.5)$$

*Proof.* We have for  $u, v \in L^\infty([0, T])$ :

$$\begin{aligned} d(\xi_t(u)\xi_t(v)) &= u_t\xi_{t-}(u)\xi_{t-}(v)dM_t + v_t\xi_{t-}(v)\xi_{t-}(u)dM_t \\ &\quad + v_tu_t\xi_{t-}(v)\xi_{t-}(u)d[M, M]_t \\ &= u_t\xi_{t-}(u)\xi_{t-}(v)dM_t + v_t\xi_{t-}(v)\xi_{t-}(u)dM_t + v_tu_t\xi_t(v)\xi_t(u)\alpha_t^2 dt \\ &\quad + \phi_tu_tv_t\xi_{t-}(v)\xi_{t-}(u)dM_t \\ &= v_tu_t\xi_t(v)\xi_t(u)\alpha_t^2 dt + \xi_{t-}(v)\xi_{t-}(u)(u_t + v_t + \phi_tu_tv_t)dM_t. \end{aligned}$$

hence

$$\begin{aligned} d(e^{-\int_0^t u_s v_s \alpha_s^2 ds} \xi_t(u)\xi_t(v)) \\ = e^{-\int_0^t u_s v_s \alpha_s^2 ds} \xi_{t-}(v)\xi_{t-}(u)(u_t + v_t + \phi_tu_tv_t)dM_t, \end{aligned}$$

which shows that  $\exp(-\langle u, v \rangle_{L_\alpha^2([0, T])})\xi(u)\xi(v) = \xi(u + v + \phi uv)$ . Relation (2.2.5) follows then by comparison with (2.2.3).  $\square$

## 2.3 Change of variable formula

We recall the following change of variable formula, which follows from Prop. 2 of [É90] after addition of an absolutely continuous drift term.

**Proposition 2.3** Let  $X = (X_t)_{t \in [0, T]}$  be a  $\mathbb{R}^n$ -valued process satisfying

$$dX_t = R_t dt + K_t dM_t, \quad X_0 > 0,$$

where  $R = (R_t)_{t \in [0, T]}$  and  $K = (K_t)_{t \in [0, T]}$  are two predictable square-integrable  $\mathbb{R}^n$ -valued processes. For any function  $\mathbb{R}_+ \times \mathbb{R}^n \ni (t, x) \rightarrow f_t(x)$  in  $\mathcal{C}_b^2(\mathbb{R}_+ \times \mathbb{R}^n; \mathbb{R})$  we have

$$f_t(X_t) = f_0(X_0) + \int_0^t L_s f_s(X_s) dM_s + \int_0^t U_s f_s(X_s) ds + \int_0^t \frac{\partial f_s}{\partial s}(X_s) ds, \quad (2.3.1)$$

where

$$L_s f_s(X_s) = i_s \langle K_s, \nabla f_s(X_s) \rangle + \frac{j_s}{\phi_s} (f_s(X_{s-} + \phi_s K_{s-}) - f_s(X_{s-})),$$

and

$$U_s f_s(X_s) = R_s \nabla f_s(X_s)$$



$$\begin{aligned}
& +\alpha_s^2 \left( \frac{1}{2} i_s \langle \text{Hess} f_s(X_s), K_s \otimes K_s \rangle \right. \\
& \left. + \frac{j_s}{\phi_s^2} (f_s(X_{s^-} + \phi_s K_{s^-}) - f_s(X_{s^-}) - \phi_s \langle K_s, \nabla f_s(X_s) \rangle) \right),
\end{aligned}$$

with the convention  $0/0 = 0$ .

## 2.4 Girsanov theorem

The Girsanov theorem holds in a particular form when  $(M_t)_{t \in [0, T]}$  is the solution of a structure equation (1.0.1). Let  $(\psi_t)_{t \in [0, T]}$  be a bounded predictable process such that  $1 + \phi_t \psi_t > 0$ ,  $t \in [0, T]$ , let  $(l_t)_{t \in [0, T]}$  denote the solution of the equation

$$dl_t = l_{t^-} \psi_t dM_t, \quad t \in [0, T], \quad l_0 = 1,$$

and let  $Q$  be the probability defined by

$$l_t = E \left[ \frac{dQ}{dP} \mid \mathcal{F}_t \right], \quad t \in [0, T]. \quad (2.4.1)$$

**Proposition 2.4** *Under the probability  $Q$ , the process*

$$Z_t = M_t - \int_0^t \alpha_s^2 \psi_s ds, \quad t \in [0, T], \quad (2.4.2)$$

*is a local martingale which satisfies the structure equation*

$$d[Z, Z]_t = \alpha_t^2 (1 + \phi_t \psi_t) dt + \phi_t dZ_t, \quad t \in [0, T]. \quad (2.4.3)$$

*Proof.* From the Girsanov theorem,

$$dZ_t = dM_t - \frac{1}{l_{t^-}} d\langle L, M \rangle_t = dM_t - \alpha_t^2 \psi_t dt$$

is a local martingale under  $Q$ , with  $d\langle L, M \rangle_t = l_{t^-} \alpha_t^2 \psi_t dt$ , and

$$d[Z, Z]_t = d[M, M]_t = \alpha_t^2 dt + \phi_t dM_t = \alpha_t^2 (1 + \phi_t \psi_t) dt + \phi_t dZ_t.$$

□

If  $(\psi_t)_{t \in [0, T]}$  and  $(\phi_t)_{t \in [0, T]}$  are deterministic, then  $(Z_t)_{t \in [0, T]}$  has the chaos representation property under  $Q$ , since (2.4.3) is a deterministic structure equation.

### 3 Computations of predictable representations

#### 3.1 Clark formula

The Clark-Ocone formula, cf. [Cla70], [KO91] is a consequence of the chaos representation property for  $(M_t)_{t \in [0, T]}$ , and states that any  $F \in \text{Dom}(D) \subset L^2(\Omega, \mathcal{F}_T, P)$  has a representation

$$F = E[F] + \int_0^T E[D_t F \mid \mathcal{F}_t] dM_t. \quad (3.1.1)$$

It can be proved as follows:

$$\begin{aligned} F &= E[F] + \sum_{n=1}^{\infty} n! \int_0^T \int_0^{t_n} \cdots \int_0^{t_2} f_n(t_1, \dots, t_n) dM_{t_1} \cdots dM_{t_n} \\ &= E[F] + \sum_{n=1}^{\infty} n \int_0^T I_{n-1}(f_n(*, t) 1_{\{*\leq t\}}) dM_t \\ &= E[F] + \int_0^T E[D_t F \mid \mathcal{F}_t] dM_t. \end{aligned}$$

Although  $D : L^2(\Omega, \mathcal{F}_T, P) \longrightarrow L^2(\Omega \times [0, T], dP \times \alpha_t^2 dt)$  is unbounded, the representation formula (3.1.1) can be extended to  $F \in L^2(\Omega, \mathcal{F}_T, P)$ .

**Proposition 3.1** *The operator  $F \mapsto E[D_t F \mid \mathcal{F}_t]$  taking its values in the space of square-integrable adapted processes has a continuous extension from  $\text{Dom}(D)$  to  $L^2(\Omega, \mathcal{F}_T, P)$ .*

*Proof.* We use the bound:

$$\begin{aligned} \|E[D_t F \mid \mathcal{F}_t]\|_{L^2(\Omega \times [0, T])}^2 &= \|E[F] - F\|_{L^2(\Omega)}^2 = \text{var}(F) \\ &\leq \|F\|_{L^2(\Omega)}^2, \quad F \in \text{Dom}(D). \end{aligned}$$

□

Instead of the adapted projection  $(E[D_t F \mid \mathcal{F}_t])_{t \in [0, T]}$  one may also use the predictable projection  $(E[D_t F \mid \mathcal{F}_{t-}])_{t \in [0, T]}$ . Here this leads to the same representation since both processes coincide in  $L^2(\Omega \times [0, T], dP \times \alpha_t^2 dt)$ .

#### 3.2 Deterministic structure equation

In this subsection we consider the case where  $(\phi_t)_{t \in [0, T]} \in L^\infty([0, T])$  is a deterministic function. In this case, the probabilistic interpretation of  $D_t$

is known and  $D_t F$  is explicitly computable. We define the operator  $D^B : \mathcal{S} \rightarrow L^2(\Omega \times \mathbb{R}_+, dP \times \alpha_t^2 dt)$  on  $\mathcal{S}$  as

$$\langle D^B F, u \rangle_{L^2_\alpha([0, T])} = \frac{d}{d\varepsilon} F \left( \omega(\cdot) + \varepsilon \int_0^\cdot i_s u_s ds \right) \Big|_{\varepsilon=0}, \quad F \in \mathcal{S}.$$

We have for  $F = \xi(u)$  and  $g \in L^2_\alpha([0, T])$ :

$$\begin{aligned} \langle D^B F, g \rangle_{L^2_\alpha([0, T])} &= \frac{d}{d\varepsilon} \exp \left( \varepsilon \int_0^T g_s u_s \alpha_s i_s ds \right) \xi(u) \Big|_{\varepsilon=0} \\ &= \int_0^T g_s u_s \alpha_s i_s ds \xi(u), \end{aligned}$$

hence  $D_t^B \xi(u) = i_t u_t \xi(u)$ ,  $t \in [0, T]$ , where

$$\xi(u) = \exp \left( \int_0^T u_s dM_s - \frac{1}{2} \int_0^T u_s^2 \alpha_s^2 i_s ds \right) \prod_{s \in J_M^T} (1 + u_s \phi_s) e^{-u_s \phi_s}. \quad (3.2.1)$$

Note that the definition of  $D^B F$  by duality in  $L^2_\alpha([0, T])$  implies

$$D_t^B \int_0^T u_s \alpha_s dB_s = u_t, \quad t \in [0, T].$$

We define a linear transformation  $T_t^\phi$  of exponential vectors, and more generally of elements of  $\mathcal{S}$ , as

$$T_t^\phi \xi(u) = (1 + u_t \phi_t) \xi(u), \quad u \in L^\infty([0, T]).$$

The transformation  $T_t^\phi$  is well-defined on  $\mathcal{S}$  because  $\xi(u_1), \dots, \xi(u_n)$  are linearly independent if  $u_1, \dots, u_n$  are distinct elements of  $L^2(\mathbb{R}_+)$ . Since we have  $\Delta M_t = 0$ ,  $dt \times dP$ -a.e.,  $T_t^\phi \xi(u)$  coincides  $dt \times dP$ -a.e. with the value at time  $T$  of the solution of the equation

$$Z_s^t = 1 + \int_0^s Z_{\tau-}^t u_\tau dM_\tau^t, \quad s \in [0, T], \quad (3.2.2)$$

where  $(M_s^t)_{s \in [0, T]}$  is defined as

$$M_s^t = M_s + \phi_t 1_{[t, T]}(s), \quad s \in [0, T].$$

In order to see this we check that  $Z_s^t = \xi_s(u)$ ,  $s < t$ ,

$$Z_t^t = (1 + \phi_t u_t) Z_{t-}^t = (1 + \phi_t u_t) \xi_{t-}(u) = (1 + \phi_t u_t) \xi_t(u),$$

(since  $\xi_{t-}(u) = \xi_t(u)$  a.s. for fixed  $t$ ), and for  $s > t$ :

$$\begin{aligned} Z_s^t &= Z_t^t + \int_t^s Z_{\tau-}^t u_\tau dM_\tau \\ &= (1 + \phi_t u_t) \xi_t(u) + \int_t^s Z_{\tau-}^t u_\tau dM_\tau, \end{aligned}$$

hence

$$\frac{Z_s^t}{1 + \phi_t u_t} = \xi_t(u) + \int_t^s \frac{Z_{\tau-}^t}{1 + \phi_t u_t} u_\tau dM_\tau, \quad s > t,$$

which implies from (2.2.3):

$$\frac{Z_s^t}{1 + \phi_t u_t} = \xi_s(u), \quad s > t,$$

and  $Z_T^t = (1 + \phi_t u_t) \xi(u) = T_t^\phi \xi(u)$ ,  $P$ -a.s.,  $t \in [0, T]$ .

In other terms,  $T_t^\phi F$ ,  $F \in \mathcal{S}$ , can be interpreted as the evaluation of  $F$  on the trajectories of  $(M_s)_{s \in [0, T]}$  perturbed by addition of a jump of height  $\phi_t$  at time  $t$ .

**Proposition 3.2** *The transformation  $T_t^\phi$  is multiplicative, i.e.*

$$T_t^\phi(FG) = (T_t^\phi F)(T_t^\phi G), \quad F, G \in \mathcal{S}.$$

Moreover we have

$$D_t F = D_t^B F + \frac{j_t}{\phi_t} (T_t^\phi F - F), \quad t \in [0, T], \quad F \in \mathcal{S}, \quad (3.2.3)$$

and

$$D_t(FG) = F D_t G + G D_t F + \phi_t D_t F D_t G, \quad t \in [0, T], \quad F, G \in \mathcal{S}. \quad (3.2.4)$$

*Proof.* For the multiplicativity we note that

$$\begin{aligned} T_t^\phi(\xi(u)\xi(v)) &= \exp(\langle u, v \rangle_{L_\alpha^2([0, T])}) T_t^\phi \xi(u + v + \phi uv) \\ &= \exp(\langle u, v \rangle_{L_\alpha^2([0, T])}) (1 + \phi_t(u_t + v_t + \phi_t u_t v_t)) \xi(u + v + \phi uv) \\ &= (1 + \phi_t u_t)(1 + \phi_t v_t) \xi(u) \xi(v) \\ &= T_t^\phi \xi(u) T_t^\phi \xi(v). \end{aligned}$$

When  $\phi_t = 0$  we have  $D_t^B F = i_t u_t \xi(u) = i_t D_t F$ , hence

$$\begin{aligned} D_t \xi(u) &= i_t D_t \xi(u) + j_t D_t \xi(u) \\ &= i_t u_t \xi(u) + j_t u_t \xi(u) \end{aligned}$$

$$= D_t^B \xi(u) + \frac{\dot{J}_t}{\phi_t} (T_t^\phi \xi(u) - \xi(u)) \quad t \in [0, T].$$

Concerning the product rule we have from Lemma 2.2:

$$\begin{aligned} D_t(\xi(u)\xi(v)) &= \exp\left(\int_0^T u_s v_s \alpha_s^2 ds\right) D_t \xi(u + v + \phi uv) \\ &= \exp\left(\int_0^T u_s v_s \alpha_s^2 ds\right) (u_t + v_t + \phi_t u_t v_t) \xi(u + v + \phi uv) \\ &= (u_t + v_t + \phi_t u_t v_t) \xi(u) \xi(v) \\ &= \xi(u) D_t \xi(v) + \xi(v) D_t \xi(u) + \phi_t D_t \xi(u) D_t \xi(v), \end{aligned}$$

$u, v \in L^\infty([0, T])$ , see also Relation (6) of [Pri96].

□

If  $(\phi_t)_{t \in [0, T]}$  is random the probabilistic interpretation of  $D$  is unknown, however we have the product rule

$$E[D_t(FG) | \mathcal{F}_t] = E[FD_tG | \mathcal{F}_t] + E[GD_tF | \mathcal{F}_t] + \phi_t E[D_tFD_tG | \mathcal{F}_t], \quad (3.2.5)$$

$F, G \in \mathcal{S}$ ,  $t \in [0, T]$ . cf. Prop. 5 of [PSV00].

### 3.3 Markovian case

This section presents a representation method which is based on the Itô formula and the Markov property, see also [Pro01] in the continuous case. Let  $(X_t)_{t \in [0, T]}$  be a  $\mathbb{R}^n$ -valued Markov (not necessarily time homogeneous) process defined on  $\Omega$ , satisfying a change of variable formula of the form

$$f(X_t) = f(X_0) + \int_0^t L_s f(X_s) dM_s + \int_0^t U_s f(X_s) ds, \quad t \in [0, T], \quad (3.3.1)$$

where  $L_s, U_s$  are operators defined on  $\mathcal{C}^2$  functions. We assume that the semi-group  $(P_{s,t})_{0 \leq s \leq t \leq T}$  associated to  $(X_t)_{t \in [0, T]}$ , i.e.

$$P_{s,t} f(X_s) = E[f(X_t) | \mathcal{F}_s] = E[f(X_t) | X_s], \quad 0 \leq s \leq t \leq T,$$

acts on  $\mathcal{C}_b^2(\mathbb{R}^n)$  functions, with  $P_{s,t} \circ P_{t,u} = P_{s,u}$ ,  $0 \leq s \leq t \leq u \leq T$ .

Although the probabilistic interpretation of  $D$  is not known when  $(\phi_t)_{t \in [0, T]}$  is random, it is still possible to compute the explicit predictable representation of  $f(X_T)$  using the Itô formula and the Markov property.

**Lemma 3.3** *Let  $f \in \mathcal{C}_b^2(\mathbb{R}^n)$ . We have*

$$E[D_t f(X_T) | \mathcal{F}_t] = (L_t(P_{t,T} f))(X_t), \quad t \in [0, T]. \quad (3.3.2)$$

*Proof.* We apply the change of variable formula (3.3.1) to  $t \mapsto P_{t,T}f(X_t) = E[f(X_T) \mid \mathcal{F}_t]$ , since  $P_{t,T}f$  is  $\mathcal{C}^2$ . Using the fact that the finite variation term vanishes since  $t \mapsto P_{t,T}f(X_t)$  is a martingale, (see e.g. Cor. 1, p. 64 of [Pro90]), we obtain:

$$P_{t,T}f(X_t) = P_{0,T}f(X_0) + \int_0^t (L_s(P_{s,T}f))(X_s) dM_s, \quad t \in [0, T],$$

with  $P_{0,T}f(X_0) = E[f(X_T)]$ . Letting  $t = T$ , we obtain (3.3.2) by uniqueness of the representation (3.1.1) applied to  $F = f(X_T)$ . □

In practice, we will use Prop. 3.1 to extend  $(E[D_t f(X_T) \mid \mathcal{F}_t])_{t \in [0, T]}$  to a less regular function  $f : \mathbb{R}^n \rightarrow \mathbb{R}$ . As an example, if  $\phi_t$  is written as  $\phi_t = \varphi(t, M_t)$ , and  $dS_t = \sigma(t, S_t)dM_t + \mu(t, S_t)dt$ , we can apply Prop. 2.3, with  $(X_t)_{t \in [0, T]} = ((S_t, M_t))_{t \in [0, T]}$  and

$$\begin{aligned} L_t f(S_t, M_t) &= i_t \sigma(t, S_t) \partial_1 f(S_t, M_t) + i_t \partial_2 f(S_t, M_t) \\ &\quad + \frac{\dot{j}_t}{\varphi(t, M_t)} (f(S_t + \varphi(t, M_t) \sigma(t, S_t), M_t + \varphi(t, M_t)) - f(S_t, M_t)), \end{aligned}$$

since the eventual jump of  $(M_t)_{t \in [0, T]}$  at time  $t$  is  $\varphi(t, M_t)$ . Here,  $\partial_1$ , resp.  $\partial_2$ , denotes the partial derivative with respect to the first, resp. second, variable. Hence

$$\begin{aligned} E[D_t f(S_T, M_T) \mid \mathcal{F}_t] &= i_t \sigma(t, S_t) (\partial_1 P_{t,T}f)((S_t, M_t)) + i_t (\partial_2 P_{t,T}f)((S_t, M_t)) \\ &\quad + \frac{\dot{j}_t}{\varphi(t, M_t)} (P_{t,T}f)((S_t + \varphi(t, M_t) \sigma(t, S_t), M_t + \varphi(t, M_t))) \\ &\quad - \frac{\dot{j}_t}{\varphi(t, M_t)} (P_{t,T}f)((S_t, M_t)). \end{aligned}$$

If  $(M_t)_{t \in [0, T]}$  is an Azéma martingale ( $\phi_t = \beta M_t$ ,  $t \in [0, T]$ ), then  $i_t = 0$   $dP \times dt$  a.s.

## 4 Computations of hedging strategies

### 4.1 Market model

In this subsection we introduce the price process which will be considered in the sequel. Let  $\mu : [0, T] \rightarrow \mathbb{R}$  and  $\sigma : [0, T] \rightarrow ]0, \infty[$  be deterministic

bounded functions. Let  $(r_t)_{t \in [0, T]}$  be a deterministic non negative function which models a risk-less asset, and let  $(\psi_t)_{t \in [0, T]}$  be defined as

$$\psi_t = \frac{r_t - \mu_t}{\sigma_t \alpha_t^2}, \quad t \in [0, T].$$

We assume that  $1 + \phi_t \psi_t > 0$ ,  $t \in [0, T]$ . If  $(\phi_t)_{t \in [0, T]}$  is not deterministic this choice is still possible due to the boundedness of  $(\phi_t)_{t \in [0, T]}$  or from Th. 2.1-iii) of [DP99]. Let  $Q$  denote the probability defined by  $E \left[ \frac{dQ}{dP} \mid \mathcal{F}_t \right] = l_t$ ,  $t \in [0, T]$ , where  $dl_t = l_t - \psi_t dM_t$ ,  $t \in [0, T]$ ,  $l_0 = 1$ . From Prop. 2.4,

$$Z_t = M_t - \int_0^t \alpha_s^2 \psi_s ds, \quad t \in [0, T],$$

is a local martingale under  $Q$  with angle bracket

$d\langle Z, Z \rangle_t = \alpha_t^2 (1 + \phi_t \psi_t) dt$ . If  $(\phi_t)_{t \in [0, T]}$  is deterministic then  $(Z_t)_{t \in [0, T]}$  is a true martingale under  $Q$ . This also holds if

$$(\phi_t)_{t \in [0, T]} = (\beta M_t)_{t \in [0, T]},$$

from the boundedness of  $(M_t)_{t \in [0, T]}$ . If  $(\phi_t)_{t \in [0, T]}$  is deterministic, we may start from the data of  $(\tilde{\alpha}_t)_{t \in [0, T]}$ , take  $d[Z, Z]_t = \tilde{\alpha}_t^2 dt + \phi_t dZ_t$  and define  $(\alpha_t)_{t \in [0, T]}$  by  $\alpha_t^2 = \tilde{\alpha}_t^2 / (1 + \phi_t \psi_t)$ ,  $t \in [0, T]$ . Let the price  $(S_t)_{t \in [0, T]}$  of a risky asset satisfy the equation

$$dS_t = \mu_t S_t dt + \sigma_t S_t - dZ_t, \quad t \in [0, T], \quad S_0 = 1. \quad (4.1.1)$$

We have

$$dS_t = r_t S_t dt + \sigma_t S_t - dM_t, \quad t \in [0, T], \quad (4.1.2)$$

and under  $P$ ,  $(S_t e^{-\int_0^t r_s ds})_{t \in [0, T]}$  is a martingale, i.e. the market is arbitrage free. Let  $\eta_t$  and  $\zeta_t$  denote the number of units invested at time  $t$  in the risky and risk-less assets respectively. Thus the value  $V_t$  of the portfolio at time  $t$  is given by

$$V_t = \zeta_t A_t + \eta_t S_t, \quad t \in [0, T], \quad (4.1.3)$$

where

$$dA_t = r_t A_t dt, \quad A_0 = 1, \quad t \in [0, T]. \quad (4.1.4)$$

We assume that the portfolio is self-financing, i.e.

$$dV_t = \zeta_t dA_t + \eta_t dS_t, \quad t \in [0, T],$$

therefore

$$dV_t = r_t V_t dt + \sigma_t \eta_t S_{t-} dM_t, \quad t \in [0, T],$$

and

$$V_T e^{-\int_0^T r_t dt} = V_0 + \int_0^T \sigma_t \eta_t S_{t-} e^{-\int_0^t r_s ds} dM_t, \quad (4.1.5)$$

Suppose that we are required to find a portfolio  $(\zeta_t, \eta_t)_{t \in [0, T]}$  which leads to a given value  $V_T = F$ . By the Clark-Ocone formula,

$$F = E[F] + \int_0^T E[D_t F | \mathcal{F}_t] dM_t,$$

and comparing with (4.1.5) we obtain

$$\begin{aligned} V_0 &= e^{-\int_0^T r_s ds} E[F], \\ \eta_t &= \sigma_t^{-1} S_t^{-1} E[D_t F | \mathcal{F}_t] e^{-\int_t^T r_s ds}, \quad t \in [0, T]. \end{aligned} \quad (4.1.6)$$

Next we consider different models with explicit computations of hedging strategies.

## 4.2 European options and deterministic structure

In this section we hedge European calls using the Clark formula. Let  $(M_t)_{t \in [0, T]}$  be the martingale described in Sect. 2.2-a), with deterministic  $(\phi_t)_{t \in [0, T]}$ , i.e.  $(M_t)_{t \in [0, T]}$  is alternatively Brownian or Poisson depending on the vanishing of  $(\phi_t)_{t \in [0, T]}$ . We assume that  $1 + \sigma_t \phi_t > 0$ ,  $t \in [0, T]$ . We have

$$S_t = \exp \left( \int_0^t \sigma_s \alpha_s i_s dB_s + \int_0^t (r_s - \phi_s \lambda_s \sigma_s - \frac{1}{2} i_s \sigma_s^2 \alpha_s^2) ds \right) \prod_{k=1}^{k=N_t} (1 + \sigma_{T_k} \phi_{T_k}),$$

$0 \leq t \leq T$ , where  $(T_k)_{k \geq 1}$  denotes the jump times of  $(N_t)_{t \in \mathbb{R}_+}$ . We will denote by  $(S_{t,u}^x)_{u \in [t, T]}$  the process defined as

$$dS_{t,u}^x = r_t S_{t,u}^x du + \sigma_u S_{t,u}^x dM_u, \quad u \in [t, T], \quad S_{t,t}^x = x.$$

We have

$$\begin{aligned} S_{t,T}^x &= x \exp \left( \int_t^T \sigma_u \alpha_u i_u dB_u + \int_t^T (r_u - \phi_u \lambda_u \sigma_u - \frac{1}{2} i_u \sigma_u^2 \alpha_u^2) du \right) \\ &\quad \times \prod_{k=1+N_t}^{k=N_T} (1 + \sigma_{T_k} \phi_{T_k}), \end{aligned}$$

$0 \leq t \leq T$ , with  $S_t = S_{0,t}^T$ ,  $t \in [0, T]$ .



**Proposition 4.1** Assume that  $\phi_t \geq 0$ ,  $t \in [0, T]$ . Then for  $0 \leq t \leq T$  we have

$$E[D_t(S_T - K)^+ | \mathcal{F}_t] = E \left[ i_t \sigma_t S_{t,T}^x 1_{[K, \infty[}(S_{t,T}^x) + \frac{\dot{J}_t}{\phi_t} (\sigma_t \phi_t S_{t,T}^x - (K - S_{t,T}^x)^+) 1_{[\frac{K}{1+\sigma_t}, \infty[}(S_{t,T}^x) \right]_{x=S_t}}.$$

*Proof.* By Prop. 3.2, for any  $F \in \mathcal{S}$  we have

$$D_t F = D_t^B F + \frac{\dot{J}_t}{\phi_t} (T_t^\phi F - F), \quad t \in [0, T]. \quad (4.2.1)$$

We have  $T_t^\phi S_T = (1 + \sigma_t \phi_t) S_T$ ,  $t \in [0, T]$ , and the chain rule  $D^B f(F) = f'(F) D^B F$  holds for  $F \in \mathcal{S}$  and  $f \in \mathcal{C}_b^2(\mathbb{R})$ . Since  $\mathcal{S}$  is an algebra for deterministic  $(\phi_t)_{t \in [0, T]}$ , we may approach  $x \mapsto (x - K)^+$  by polynomials on compact intervals and proceed e.g. as in [Øks96], p. 5-13. By dominated convergence,  $(S_T - K)^+ \in \text{Dom}(D)$  and (4.2.1) becomes

$$D_t(S_T - K)^+ = i_t \sigma_t S_T 1_{[K, \infty[}(S_T) + \frac{\dot{J}_t}{\phi_t} ((1 + \sigma_t \phi_t) S_T - K)^+ - (S_T - K)^+,$$

$0 \leq t \leq T$ . The Markov property of  $(S_t)_{t \in [0, T]}$  implies

$$E[D_t^B(S_T - K)^+ | \mathcal{F}_t] = i_t \sigma_t E[S_{t,T}^x 1_{[K, \infty[}(S_{t,T}^x)]_{x=S_t}},$$

and

$$\begin{aligned} & \frac{\dot{J}_t}{\phi_t} E \left[ (T_t^\phi S_T - K)^+ - (S_T - K)^+ | \mathcal{F}_t \right] \\ &= \frac{\dot{J}_t}{\phi_t} E \left[ ((1 + \sigma_t \phi_t) S_{t,T}^x - K)^+ - (S_{t,T}^x - K)^+ \right]_{x=S_t} \\ &= \frac{\dot{J}_t}{\phi_t} E \left[ ((1 + \sigma_t \phi_t) S_{t,T}^x - K) 1_{[\frac{K}{1+\sigma_t}, \infty[}(S_{t,T}^x) \right]_{x=S_t} \\ &\quad - \frac{\dot{J}_t}{\phi_t} E \left[ (S_{t,T}^x - K)^+ 1_{[K, \infty[}(S_{t,T}^x) \right]_{x=S_t} \\ &= \frac{\dot{J}_t}{\phi_t} E \left[ \sigma_t \phi_t S_{t,T}^x 1_{[\frac{K}{1+\sigma_t}, \infty[}(S_{t,T}^x) + (S_{t,T}^x - K) 1_{[\frac{K}{1+\sigma_t}, K]}(S_{t,T}^x) \right]_{x=S_t} \\ &= \frac{\dot{J}_t}{\phi_t} E \left[ \sigma_t \phi_t S_{t,T}^x 1_{[\frac{K}{1+\sigma_t}, \infty[}(S_{t,T}^x) - (K - S_{t,T}^x)^+ 1_{[\frac{K}{1+\sigma_t}, \infty[}(S_{t,T}^x) \right]_{x=S_t} \\ &= \frac{\dot{J}_t}{\phi_t} E \left[ (\sigma_t \phi_t S_{t,T}^x - (K - S_{t,T}^x)^+) 1_{[\frac{K}{1+\sigma_t}, \infty[}(S_{t,T}^x) \right]_{x=S_t}. \end{aligned}$$

□

If  $(\phi_t)_{t \in [0, T]}$  is not constrained to be positive then

$$\begin{aligned} E[D_t(S_T - K)^+ | \mathcal{F}_t] &= i_t \sigma_t E \left[ S_{t,T}^x 1_{[K, \infty[}(S_{t,T}^x) \right]_{x=S_t} \\ &+ \frac{j_t}{\phi_t} E \left[ \sigma_t \phi_t S_{t,T}^x 1_{[\frac{K}{1+\sigma_t \phi_t}, \infty[}(S_{t,T}^x) + (S_{t,T}^x - K) 1_{[\frac{K}{1+\sigma_t \phi_t}, K]}(S_{t,T}^x) \right]_{x=S_t}, \end{aligned}$$

with the convention  $1_{[b,a]} = -1_{[a,b]}$ ,  $0 \leq a < b \leq T$ . Prop. 4.1 can also be proved using Lemma 3.3 and the Itô formula (2.3.1). In the deterministic case, the semi-group  $P_{t,T}$  can be explicitly computed. Let

$\Gamma_{t,T}^\sigma = \int_t^T i_s \alpha_s^2 \sigma_s^2 ds$  denote the variance of  $\int_t^T i_s \alpha_s \sigma_s dB_s$ ,  $t \in [0, T]$ , and let  $\Gamma_{t,T} = \int_t^T \gamma_s ds$ ,  $t \in [0, T]$ , denote the intensity of  $N_T - N_t$  under  $Q$ , where  $\gamma_t = \lambda_t(1 + \phi_t \psi_t)$ ,  $t \in [0, T]$ .

**Proposition 4.2** *We have for  $f \in \mathcal{C}_b(\mathbb{R})$*

$$\begin{aligned} P_{t,T}f(x) &= \sum_{k=0}^{\infty} \frac{e^{-\Gamma_{t,T}}}{k!} \int_{-\infty}^{\infty} \nu_{t_0} \int_{[t,T]^k} \gamma_{t_1} \cdots \gamma_{t_k} \\ &f \left( x e^{-\frac{\Gamma_{t,T}^\sigma}{2} + (\Gamma_{t,T}^\sigma)^{1/2} t_0 - \int_t^T \phi_s \gamma_s \sigma_s ds} \prod_{i=1}^k (1 + \sigma_{t_i} \phi_{t_i}) \right) dt_1 \cdots dt_k dt_0, \end{aligned}$$

where  $\nu$  denotes the standard Gaussian density.

*Proof.* We have  $P_{t,T}f(x) = E[f(S_T) | S_t = x] = E[f(S_{t,T}^x)]$ , and

$$P_{t,T}f(x) = \sum_{k=0}^{\infty} E[f(S_{t,T}^x) | N_T - N_t = k] \exp(-\Gamma_{t,T}) \frac{\Gamma_{t,T}^k}{k!}$$

$k \in \mathbb{N}$ . Since  $(N_{\Gamma_{t,s}^{-1}} - N_t)_{s \in [t, T]}$  is a standard Poisson process, conditionally to  $\{N_T - N_t = k\}$ , the  $n$  first jump times  $(T_1, \dots, T_n)$  of  $(N_s)_{s \in [t, T]}$  have the law

$$\frac{k!}{(\Gamma_{t,T})^k} 1_{\{t < t_1 < \dots < t_k < T\}} \gamma_{t_1} \cdots \gamma_{t_k} dt_1 \cdots dt_k,$$

and conditionally to  $\{N_T - N_t = k\}$ , the jump times  $(\Gamma_{t,T_1}, \dots, \Gamma_{t,T_k})$  have a uniform law on  $[0, \Gamma_{t,T}]^k$ . We then use the identity in law between  $S_{t,T}^x$  and

$$x X_{t,T} \exp \left( - \int_t^T \phi_s \lambda_s (1 + \phi_s \psi_s) \sigma_s ds \right) \prod_{k=1}^{k=N_T} (1 + \sigma_{T_k} \phi_{T_k}),$$

where

$$X_{t,T} = \exp \left( -\Gamma_{t,T}^\sigma / 2 + (\Gamma_{t,T}^\sigma)^{1/2} W \right),$$

and  $W$  a standard Gaussian random variable, independent of  $(N_t)_{t \in [0, T]}$ . This identity holds because  $(B_t)_{t \in [0, T]}$  is a standard Brownian motion, independent of  $(N_t)_{t \in [0, T]}$ . □

See Prop. 8 of [JP02] for a computation of

$$E \left[ \exp \left( - \int_0^T r_s ds \right) (S_T - K)^+ \right]$$

in terms of the classical Black-Scholes function

$$\text{BS}(x, T; r, \sigma^2; K) = E[e^{-rT} (xe^{rT - \sigma^2 T/2 + \sigma W_t} - K)^+],$$

where  $W_t$  is a centered Gaussian random variable with variance  $t$ .

### 4.3 Asian options and deterministic structure

The price at time  $t$  of such an option is

$$E \left[ e^{-\int_t^T r_s ds} \left( \frac{1}{T} \int_0^T S_u du - K \right)^+ \mid \mathcal{F}_t \right].$$

The next proposition gives us a replicating hedging strategy for Asian options in the the case of deterministic structure equation model

**Proposition 4.3** *There exists a measurable function  $\tilde{C}$  on  $\mathbb{R}_+ \times \mathbb{R}$  such that  $\tilde{C}(t, \cdot)$  is  $\mathcal{C}^1$  for all  $t \in \mathbb{R}_+$ , and*

$$S_t \tilde{C}(t, Y_t) = E \left[ \left( \frac{1}{T} \int_0^T S_u du - K \right)^+ \mid \mathcal{F}_t \right].$$

Moreover, the replicating portfolio for an Asian option with payoff

$$\left( \frac{1}{T} \int_0^T S_u du - K \right)^+$$

is given by (4.1.3) and

$$\begin{aligned} \eta_t = & \frac{1}{\sigma_t} e^{-\int_t^T r_s ds} \left[ \tilde{C}(t, Y_t) \sigma_t \right. \\ & \left. + (1 + \sigma_t \phi_t) \left( \frac{j_t}{\phi_t} \left( \tilde{C} \left( t, \frac{Y_t}{1 + \sigma_t \phi_t} \right) - \tilde{C}(t, Y_t) \right) - i_t \sigma_t Y_t \partial_2 \tilde{C}(t, Y_t) \right) \right]. \end{aligned} \quad (4.3.1)$$

*Proof.* Following [LL96], p. 91, we define the auxiliary process

$$Y_t = \frac{1}{S_t} \left( \frac{1}{T} \int_0^t S_u du - K \right), \quad t \in [0, T].$$

With this notation, the price of the Asian option at time  $t$  becomes

$$E \left[ e^{-\int_t^T r_s ds} S_T (Y_T)^+ \mid \mathcal{F}_t \right].$$

For  $0 \leq s \leq t \leq T$ , we have

$$d(S_t Y_t) = \frac{1}{T} d \left( \int_0^t S_u du - K \right) = \frac{S_t}{T} dt,$$

hence

$$\frac{S_t Y_t}{S_s} = Y_s + \frac{1}{T} \int_s^t \frac{S_u}{S_s} du.$$

Let  $H \in \mathcal{C}_b^2(\mathbb{R})$ . We have

$$\begin{aligned} E[H(S_T Y_T) | \mathcal{F}_t] &= E \left[ H \left( S_t Y_t + \frac{1}{T} \int_t^T S_u du \right) | \mathcal{F}_t \right] \\ &= E \left[ H \left( xy + \frac{x}{T} \int_t^T \frac{S_u}{S_t} du \right) \right]_{y=Y_t, x=S_t}. \end{aligned}$$

Let  $C \in \mathcal{C}_b^2(\mathbb{R}_+ \times \mathbb{R}^2)$  be defined as

$$C(t, x, y) = E \left[ H \left( xy + \frac{x}{T} \int_t^T \frac{S_u}{S_t} du \right) \right],$$

i.e.

$$C(t, S_t, Y_t) = E[H(S_T Y_T) | \mathcal{F}_t].$$

When  $H(x) = x^+$ , since for any  $t \in [0, T]$ ,  $S_t$  is positive and  $\mathcal{F}_t$ -measurable, and  $S_u/S_t$  is independent of  $\mathcal{F}_t$ ,  $u \geq t$ , we have:

$$\begin{aligned} E[H(S_T Y_T) | \mathcal{F}_t] &= E[S_T (Y_T)^+ | \mathcal{F}_t] \\ &= S_t E \left[ \left( \frac{S_T}{S_t} Y_T \right)^+ | \mathcal{F}_t \right] \\ &= S_t E \left[ \left( Y_t + \frac{1}{T} \int_t^T \frac{S_u}{S_t} du \right)^+ | \mathcal{F}_t \right] \\ &= S_t E \left[ \left( y + \frac{1}{T} \int_t^T \frac{S_u}{S_t} du \right)^+ \right]_{y=Y_t} \\ &= S_t \tilde{C}(t, Y_t), \end{aligned}$$

with

$$\tilde{C}(t, y) = E \left[ \left( y + \frac{1}{T} \int_t^T \frac{S_u}{S_t} du \right)^+ \right].$$

We now proceed as in [Bel99], which deals with the sum of a Brownian motion and a Poisson process. From the expression of  $1/S_t$  we have

$$d \left( \frac{1}{S_t} \right) = \frac{1}{S_{t-}} \left[ \left( -r_t + \frac{\alpha_t^2 \sigma_t^2}{1 + \sigma_t \phi_t} \right) dt - \frac{\sigma_t}{1 + \sigma_t \phi_t} dM_t \right],$$

hence by (2.3.1)

$$dY_t = Y_t \left( -r_t + \frac{\alpha_t^2 \sigma_t^2}{1 + \sigma_t \phi_t} \right) dt + \frac{1}{T} dt - \frac{Y_t \sigma_t}{1 + \sigma_t \phi_t} dM_t.$$

Applying Lemma. 3.3 we get

$$\begin{aligned} E [D_t H(S_T Y_T) | \mathcal{F}_t] &= L_t C(t, S_t, Y_t) \\ &= i_t \left( \sigma_t S_t \partial_2 C(t, S_t, Y_t) - \frac{Y_t \sigma_t}{1 + \sigma_t \phi_t} \partial_3 C(t, S_t, Y_t) \right) \\ &\quad + \frac{j_t}{\phi_t} \left( C \left( t, S_{t-} + \sigma_t S_{t-}, Y_{t-} - \frac{Y_t \sigma_t}{1 + \sigma_t \phi_t} \right) - C(t, S_{t-}, Y_{t-}) \right). \end{aligned} \quad (4.3.2)$$

Given a family  $(H_n)_{n \in \mathbb{N}}$  of  $\mathcal{C}_b^2$  functions, such that  $|H_n(x)| \leq x^+$  and  $|H'_n(x)| \leq 2$ ,  $x \in \mathbb{R}$ ,  $n \in \mathbb{N}$ , and converging pointwise to  $x \rightarrow x^+$ , by dominated convergence (4.3.2) holds for  $C(t, x, y) = x \tilde{C}(t, y)$  and we obtain:

$$\begin{aligned} E \left[ D_t \left( \frac{1}{T} \int_0^T S_u du - K \right)^+ | \mathcal{F}_t \right] \\ &= i_t \tilde{C}(t, Y_t) \sigma_t S_t \\ &\quad + S_t \left( \frac{j_t}{\phi_t} \left( \tilde{C} \left( t, \frac{Y_t}{1 + \sigma_t \phi_t} \right) - \tilde{C}(t, Y_t) \right) - i_t \sigma_t Y_t \partial_2 \tilde{C}(t, Y_t) \right) \\ &\quad + S_t \sigma_t \phi_t \left( \frac{j_t}{\phi_t} \left( \tilde{C} \left( t, \frac{Y_t}{1 + \sigma_t \phi_t} \right) - \tilde{C}(t, Y_t) \right) - i_t \sigma_t Y_t \partial_2 \tilde{C}(t, Y_t) \right). \end{aligned}$$

□

As a particular case we consider the Brownian motion model, i.e.  $\phi_t = 0$ , for all  $t \in [0, T]$ , so  $i_t = 1$ ,  $j_t = 0$  for all  $t \in [0, T]$ , and we are in the Brownian motion model. In this case we have

$$\begin{aligned} \eta_t &= e^{-\int_t^T r_s ds} \left( -Y_t \partial_2 \tilde{C}(t, Y_t) + \tilde{C}(t, Y_t) \right) \\ &= e^{-\int_t^T r_s ds} \left( S_t \frac{\partial}{\partial x} \tilde{C} \left( t, \frac{1}{x} \left( \frac{1}{T} \int_0^t S_u du - K \right) \right) \Big|_{x=S_t} + \tilde{C}(t, Y_t) \right) \\ &= \frac{\partial}{\partial x} \left( x e^{-\int_t^T r_s ds} \tilde{C} \left( t, \frac{1}{x} \left( \frac{1}{T} \int_0^t S_u du - K \right) \right) \right) \Big|_{x=S_t}, \quad t \in [0, T], \end{aligned}$$

which can be denoted informally as a partial derivative with respect to  $S_t$ .

## 4.4 European options and Azéma martingales

Let  $-2 \leq \beta < 0$ , and let  $(M_t)_{t \in [0, T]}$  be the unique solution of the structure equation

$$d[M, M]_t = dt + \beta M_t - dM_t, \quad t \in [0, T]. \quad (4.4.1)$$

This process has the chaos representation property, hence the results of Sect. 3 apply in this case. This allows to obtain an explicit hedging formula for the model of [DP99]. We use the convention  $1_{[b, a]} = -1_{[a, b]}$ ,  $0 \leq a < b \leq T$ .

**Proposition 4.4** *We have*

$$\begin{aligned} E [D_t(S_T - K)^+ | \mathcal{F}_t] \\ &= \frac{1}{\beta M_t} E \left[ \sigma_t \beta (y + M_T - M_t) S_{t, T}^x 1_{[\frac{K}{1 + \sigma_t \beta (y + M_T - M_t)}, \infty[}(S_{t, T}^x) \right. \\ &\quad \left. + (S_{t, T}^x - K) 1_{[\frac{K}{1 + \sigma_t \beta (y + M_T - M_t)}, K]}(S_{t, T}^x) \right]_{x=S_t}^{y=M_t}. \end{aligned}$$

*Proof.* Let  $(X_t)_{t \in [0, T]} = ((S_t, M_t))_{t \in [0, T]}$ ,  $(R_t)_{t \in [0, T]} = ((r_t S_t, 0))_{t \in [0, T]}$ ,  $(K_t)_{t \in [0, T]} = ((\sigma_t S_t, 1))_{t \in [0, T]}$ , and  $X_0 = (1, 0)$ . By Lemma 3.3, for  $f \in \mathcal{C}_b^2(\mathbb{R}^2)$  we have

$$\begin{aligned} E[D_t f(X_T) | \mathcal{F}_t] &= (L_t(P_{t, T} f))(X_t) \\ &= \frac{1}{\beta M_t} ((P_{t, T} f)(X_t + \beta M_t K_t) - (P_{t, T} f)(X_t)) \\ &= \frac{1}{\beta M_t} ((P_{t, T} f)((1 + \beta M_t \sigma_t) S_t, (1 + \beta) M_t) - (P_{t, T} f)(S_t, M_t)) \\ &= \frac{1}{\beta M_t} E [f((1 + \sigma_t \beta (y + M_T - M_t)) S_{t, T}^x, \\ &\quad (1 + \beta (y + M_T - M_t)))]_{x=S_t}^{y=M_t} \\ &\quad - \frac{1}{\beta M_t} E [f(S_{t, T}^x, y + M_T - M_t)]_{x=S_t}^{y=M_t}. \end{aligned}$$

In particular if  $f(x, y) = f(x)$  depends only on the first variable we have

$$\begin{aligned} E [D_t f(S_T) | \mathcal{F}_t] \\ &= \frac{1}{\beta M_t} E [f((1 + \sigma_t \beta (y + M_T - M_t)) S_{t, T}^x) - f(S_{t, T}^x)]_{x=S_t}^{y=M_t}. \end{aligned}$$

Approaching the function  $x \mapsto (x - K)^+$  with a sequence  $(f_n)_{n \in \mathbb{N}}$  of  $\mathcal{C}_b^2$  functions converging pointwise with  $|f_n(x)| \leq (x - K)^+$ , and  $|f'_n(x)| \leq 2$ ,  $x \in \mathbb{R}$ ,  $n \in \mathbb{N}$ , we obtain

$$E [D_t(S_T - K)^+ | \mathcal{F}_t]$$

$$\begin{aligned}
&= \frac{1}{\beta M_t} E \left[ \left( (1 + \sigma_t \beta (y + M_T - M_t)) S_{t,T}^x - K \right)^+ - (S_{t,T}^x - K)^+ \right]_{x=S_t}^{y=M_t} \\
&= \frac{1}{\beta M_t} E \left[ \left( (1 + \sigma_t \beta (y + M_T - M_t)) S_{t,T}^x - K \right) 1_{\left[ \frac{K}{1 + \sigma_t \beta (y + M_T - M_t)}, \infty \right]}(S_{t,T}^x) \right. \\
&\quad \left. - (S_{t,T}^x - K) 1_{[K, \infty]}(S_{t,T}^x) \right]_{x=S_t}^{y=M_t} \\
&= \frac{1}{\beta M_t} E \left[ \sigma_t \beta (y + M_T - M_t) S_{t,T}^x 1_{\left[ \frac{K}{1 + \sigma_t \beta (y + M_T - M_t)}, \infty \right]}(S_{t,T}^x) \right. \\
&\quad \left. + (S_{t,T}^x - K) 1_{\left[ \frac{K}{1 + \sigma_t \beta (y + M_T - M_t)}, K \right]}(S_{t,T}^x) \right]_{x=S_t}^{y=M_t}.
\end{aligned}$$

□

## 4.5 Lookback options

Hedging strategies for Lookback options have been computed in [Ber98] using the Clark-Ocone formula. In this section we show that classical martingale methods also apply in this case. We assume that

$(M_t)_{t \in [0, T]} = (B_t)_{t \in [0, T]}$  is a standard Brownian motion, i.e.  $\alpha_t = 1$  and  $\phi_t = 0$  for every  $t \in [0, T]$ , and take  $r_t = r \geq 0$  and  $\sigma_t = \sigma \geq 0$  for every  $t \in [0, T]$ . Under the risk-free probability  $P$  the asset price  $(S_t)_{t \in [0, T]}$  has the dynamic

$$dS_t = rS_t dt + \sigma S_t dB_t, \quad t \in [0, T],$$

so (4.1.5) becomes

$$V_T e^{-rT} = V_0 + \int_0^T \sigma \eta_t S_t e^{-rt} dB_t, \quad t \in [0, T].$$

Let  $m_s^t = \inf_{u \in [s, t]} S_u$ ,  $M_s^t = \sup_{u \in [s, t]} S_u$ ,  $0 \leq s \leq t \leq T$ , and let  $\mathcal{M}_s^t$  be either  $m_s^t$  or  $M_s^t$ . In the Lookback option case the payoff  $H(S_T, \mathcal{M}_0^T)$  depends not only on the price of the underlying asset at maturity but also it depends on all prices of the underlying asset over the period which starts from the initial time and ends at maturity, and let  $\text{Look}_t$  be the price of the Lookback option given by

$$\text{Look}_t = e^{-r(T-t)} E[H(S_T, \mathcal{M}_0^T) \mid \mathcal{F}_t], \quad H \in \mathcal{C}_b^2(\mathbb{R}^2), \quad t \in [0, T].$$

**Proposition 4.5** *There exists a  $\mathcal{C}^1$  function  $f$  such that*

$$f(S_t, \mathcal{M}_0^t, t) = E[H(S_T, \mathcal{M}_0^T) \mid \mathcal{F}_t], \quad 0 \leq t \leq T.$$

*The replicating portfolio of a Lookback option with payoff  $H(S_T, \mathcal{M}_0^T)$ , and price  $\text{Look}_t = f(S_t, \mathcal{M}_0^t, t)$  at time  $t$ , is given by (4.1.3), and*

$$\eta_t = e^{-r(T-t)} \partial_1 f(S_t, \mathcal{M}_0^t, t), \quad t \in [0, T]. \quad (4.5.1)$$

*Proof.* It suffices to deal with the case  $\mathcal{M}_s^t = m_s^t$ . The existence of  $f$  follows from the Markov property, more precisely

$$f(x, y, t) = E [H (S_{t,T}^x, y \wedge \mathcal{M}_t^T)].$$

Applying the change of variable formula, for  $t \in [0, T]$ , we have

$$\begin{aligned} df(S_t, \mathcal{M}_0^t, t) &= \left[ \partial_3 f + r S_t \partial_1 f + \frac{1}{2} \sigma^2 S_t^2 \partial_1^2 f \right] (S_t, \mathcal{M}_0^t, t) dt \\ &\quad + \partial_2 f(S_t, \mathcal{M}_0^t, t) d\mathcal{M}_0^t + \sigma S_t \partial_1 f(S_t, \mathcal{M}_0^t, t) dB_t. \end{aligned}$$

Since  $(E[H(S_T, \mathcal{M}_0^T) | \mathcal{F}_t])_{t \in [0, T]}$  is a  $P$ -martingale and  $(\mathcal{M}_0^t)_{t \in [0, T]}$  has finite variation (it is in fact a decreasing process), we have:

$$df(S_t, \mathcal{M}_0^t, t) = \sigma S_t \partial_1 f(S_t, \mathcal{M}_0^t, t) dB_t \quad t \in [0, T].$$

Then

$$e^{-rT} F = e^{-rT} E[F] + \int_0^T e^{-rt} \sigma S_t \frac{\partial}{\partial x} f(x, \mathcal{M}_0^t, t)|_{x=S_t} dB_t \quad t \in [0, T],$$

which shows (4.5.1). □

It is stated in Bermin [Ber98] that we should have

$$\int_0^t \partial_2 f(S_s, \mathcal{M}_0^s, s) d\mathcal{M}_0^s = 0, \quad (4.5.2)$$

for the delta hedging method to work. We showed in Prop. 4.5 that the delta hedging approach can be applied without having to verify (4.5.2), since  $(\mathcal{M}_0^t)_{t \in [0, T]}$  is a monotone process with finite variation.

Relation (4.5.1) can be written informally as

$$\eta_t = \frac{\partial}{\partial S_t} \text{Look}_t, \quad t \in [0, T].$$

Let

$$d_t^T(y) = \frac{\log \frac{S_t}{y} + (r + \frac{1}{2} \sigma^2)(T - t)}{\sigma \sqrt{T - t}}, \quad \text{and} \quad \mathcal{N}(y) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^y e^{-\frac{1}{2} u^2} du.$$

A standard Lookback call option is the right to buy the underlying asset at the historically lowest price. In this case the strike is  $m_0^T$  and the payoff is

$$G = S_T - m_0^T.$$



From [DJ98], Prop. 4, p. 271, the price  $\text{Look}_t$  at time  $t$  is given by

$$\begin{aligned}
\text{Look}_t &= E_Q[e^{-r(T-t)}(S_T - m_0^T) \mid \mathcal{F}_t] & (4.5.3) \\
&= S_t \mathcal{N}(d_t^T(m_0^t)) - e^{-r(T-t)} m_0^t \mathcal{N}\left(d_t^T(m_0^t) - \sigma\sqrt{T-t}\right) \\
&\quad + e^{-r(T-t)} \frac{S_t \sigma^2}{2r} \left[ \left(\frac{S_t}{m_0^t}\right)^{-\frac{2r}{\sigma^2}} \mathcal{N}\left(-d_t^T(m_0^t) + \frac{2r\sqrt{T-t}}{\sigma}\right) \right. \\
&\quad \left. - e^{r(T-t)} \mathcal{N}(-d_t^T(m_0^t)) \right].
\end{aligned}$$

In the following proposition we recover the result of [Ber98], § 2.6.1, p. 29, using the delta hedging approach instead of the Clark formula, as an application of Prop. 4.5.

**Proposition 4.6** *The hedging strategy for a standard Lookback call option is given by*

$$\begin{aligned}
\eta_t &= \mathcal{N}(d_t^T(m_0^t)) - \frac{\sigma^2}{2r} \mathcal{N}(-d_t^T(m_0^t)) & (4.5.4) \\
&\quad + e^{-r(T-t)} \left(\frac{S_t}{m_0^t}\right)^{\frac{-2r}{\sigma^2}} \left(\frac{\sigma^2}{2r} - 1\right) \mathcal{N}\left(-d_t^T(m_0^t) + \frac{2r\sqrt{T-t}}{\sigma}\right).
\end{aligned}$$

*Proof.* We need to compute the following derivatives:

$$\begin{aligned}
&\frac{\partial}{\partial S_t} (\mathcal{N}(d_t^T(m_0^t))) \\
&= \frac{\partial}{\partial S_t} \left( \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{(d_t^T(m_0^t))} e^{-\frac{1}{2}u^2} du \right) \\
&= \frac{1}{\sqrt{2\pi}} \left[ \frac{\partial}{\partial S_t} (d_t^T(m_0^t)) \right] \exp\left(-\frac{1}{2}(d_t^T(m_0^t))^2\right) \\
&= \frac{1}{\sqrt{2\pi}} \left[ \frac{\partial}{\partial S_t} \left( \frac{\log \frac{S_t}{m_0^t} + (r + \frac{1}{2}\sigma^2)(T-t)}{\sigma\sqrt{T-t}} \right) \right] \exp\left(-\frac{1}{2}(d_t^T(m_0^t))^2\right) \\
&= \frac{1}{S_t \sigma \sqrt{2\pi(T-t)}} \exp\left(-\frac{1}{2}(d_t^T(m_0^t))^2\right),
\end{aligned}$$

and

$$\begin{aligned}
&\frac{\partial}{\partial S_t} \left( \mathcal{N}\left(d_t^T(m_0^t) - \sigma\sqrt{T-t}\right) \right) \\
&= \frac{1}{S_t \sigma \sqrt{2\pi(T-t)}} \exp\left(-\frac{1}{2}\left(d_t^T(m_0^t) - \sigma\sqrt{T-t}\right)^2\right).
\end{aligned}$$

Similarly we have

$$\frac{\partial}{\partial S_t} \left( \mathcal{N}\left(-d_t^T(m_0^t) + \frac{2r\sqrt{T-t}}{\sigma}\right) \right)$$

$$= -\frac{1}{S_t \sigma \sqrt{2\pi(T-t)}} \exp\left(-\frac{1}{2}\left(-d_t^T(m_0^t) + \frac{2r\sqrt{T-t}}{\sigma}\right)^2\right),$$

and

$$\frac{\partial}{\partial S_t} (\mathcal{N}(-d_t^T(m_0^t))) = -\frac{1}{S_t \sigma \sqrt{2\pi(T-t)}} \exp\left(-\frac{1}{2}(d_t^T(m_0^t))^2\right).$$

Finally,

$$\frac{\partial}{\partial S_t} \left(\frac{S_t}{m_0^t}\right)^{-\frac{2r}{\sigma^2}} = \frac{-2r}{m_0^t \sigma^2} \left(\frac{S_t}{m_0^t}\right)^{-\frac{2r}{\sigma^2}-1}.$$

The above expressions can be combined to compute the derivative of  $\text{Look}_t$  in (4.5.3), and to obtain:

$$\begin{aligned} \eta_t &= \mathcal{N}(d_t^T(m_0^t)) + \frac{1}{\sigma \sqrt{2\pi(T-t)}} \exp\left(-\frac{1}{2}(d_t^T(m_0^t))^2\right) \\ &\quad - e^{-r(T-t)} m_0^t \frac{1}{S_t \sigma \sqrt{2\pi(T-t)}} \exp\left(-\frac{1}{2}\left(d_t^T(m_0^t) - \sigma\sqrt{T-t}\right)^2\right) \\ &\quad + e^{-r(T-t)} \frac{\sigma^2}{2r} \left[ \left(\frac{S_t}{m_0^t}\right)^{-\frac{2r}{\sigma^2}} \mathcal{N}\left(-d_t^T(m_0^t) + \frac{2r\sqrt{T-t}}{\sigma}\right) \right. \\ &\quad \quad \left. - e^{r(T-t)} \mathcal{N}(-d_t^T(m_0^t)) \right] \\ &\quad + e^{-r(T-t)} \frac{S_t \sigma^2}{2r} \left[ \frac{-2r}{m_0^t \sigma^2} \left(\frac{S_t}{m_0^t}\right)^{-\frac{2r}{\sigma^2}-1} \mathcal{N}\left(-d_t^T(m_0^t) + \frac{2r\sqrt{T-t}}{\sigma}\right) \right. \\ &\quad \quad \left. - \left(\frac{S_t}{m_0^t}\right)^{-\frac{2r}{\sigma^2}} \frac{1}{S_t \sigma \sqrt{2\pi(T-t)}} \exp\left(-\frac{1}{2}\left(-d_t^T(m_0^t) + \frac{2r\sqrt{T-t}}{\sigma}\right)^2\right) \right. \\ &\quad \quad \left. + e^{r(T-t)} \frac{1}{S_t \sigma \sqrt{2\pi(T-t)}} \exp\left(-\frac{1}{2}(d_t^T(m_0^t))^2\right) \right] \\ &= \mathcal{N}\left(-d_t^T(m_0^t) + \frac{2r\sqrt{T-t}}{\sigma}\right) \left[ e^{-r(T-t)} \frac{\sigma^2}{2r} \left(\frac{S_t}{m_0^t}\right)^{-\frac{2r}{\sigma^2}} + e^{-r(T-t)} \frac{S_t \sigma^2}{2r} \right. \\ &\quad \quad \left. \times \frac{-2r}{m_0^t \sigma^2} \left(\frac{S_t}{m_0^t}\right)^{-\frac{2r}{\sigma^2}-1} \right] + \mathcal{N}(d_t^T(m_0^t)) \\ &\quad - \mathcal{N}(-d_t^T(m_0^t)) \left[ -e^{-r(T-t)} \frac{\sigma^2}{2r} \times e^{r(T-t)} \right] \\ &\quad + \frac{1}{\sigma \sqrt{2\pi(T-t)}} \left\{ \exp\left(-\frac{1}{2}(d_t^T(m_0^t))^2\right) \left[ 1 + e^{-r(T-t)} \frac{S_t \sigma^2}{2r} \times \frac{e^{r(T-t)}}{S_t} \right] \right. \\ &\quad \quad \left. - e^{-r(T-t)} \left[ \frac{m_0^t}{S_t} \exp\left(-\frac{1}{2}\left(d_t^T(m_0^t) - \sigma\sqrt{T-t}\right)^2\right) \right] \right\} \end{aligned}$$

$$\begin{aligned}
& + \frac{\sigma^2}{2r} \left( \frac{S_t}{m_0^t} \right)^{-\frac{2r}{\sigma^2}} \exp \left( -\frac{1}{2} \left( -d_t^T(m_0^t) + \frac{2r\sqrt{T-t}}{\sigma} \right)^2 \right) \Bigg\} \\
= & e^{-r(T-t)} \left( \frac{S_t}{m_0^t} \right)^{-\frac{2r}{\sigma^2}} \left( \frac{\sigma^2}{2r} - 1 \right) \mathcal{N} \left( -d_t^T(m_0^t) + \frac{2r\sqrt{T-t}}{\sigma} \right) \\
& + \mathcal{N} \left( d_t^T(m_0^t) \right) - \frac{\sigma^2}{2r} \mathcal{N} \left( -d_t^T(m_0^t) \right) \\
& + \frac{1}{\sigma\sqrt{2\pi(T-t)}} \left\{ \left( 1 + \frac{\sigma^2}{2r} \right) \exp \left( -\frac{1}{2} d_t^T(m_0^t)^2 \right) \right. \\
& - e^{-r(T-t)} \left[ \frac{m_0^t}{S_t} \exp \left( -\frac{1}{2} \left( d_t^T(m_0^t) - \sigma\sqrt{T-t} \right)^2 \right) \right. \\
& \left. \left. + \frac{\sigma^2}{2r} \left( \frac{S_t}{m_0^t} \right)^{-\frac{2r}{\sigma^2}} \exp \left( -\frac{1}{2} \left( -d_t^T(m_0^t) + \frac{2r\sqrt{T-t}}{\sigma} \right)^2 \right) \right] \right\}.
\end{aligned}$$

To obtain (4.5.4), it is sufficient to show that

$$\begin{aligned}
0 = & \left( 1 + \frac{\sigma^2}{2r} \right) \exp \left( -\frac{1}{2} d_t^T(m_0^t)^2 \right) \\
& - e^{-r(T-t)} \left[ \frac{m_0^t}{S_t} \exp \left( -\frac{1}{2} \left( d_t^T(m_0^t) - \sigma\sqrt{T-t} \right)^2 \right) \right. \\
& \left. + \frac{\sigma^2}{2r} \left( \frac{S_t}{m_0^t} \right)^{-\frac{2r}{\sigma^2}} \exp \left( -\frac{1}{2} \left( -d_t^T(m_0^t) + \frac{2r\sqrt{T-t}}{\sigma} \right)^2 \right) \right].
\end{aligned}$$

To see this, one can observe that

$$\begin{aligned}
& \exp \left( -\frac{1}{2} \left( d_t^T(y) - \sigma\sqrt{T-t} \right)^2 \right) \\
= & \exp \left( -\frac{1}{2} \left[ \left( d_t^T(y) \right)^2 + \sigma^2(T-t) - 2d_t^T(y)\sigma\sqrt{T-t} \right] \right) \\
= & \exp \left( -\frac{1}{2} \left( d_t^T(y) \right)^2 \right) \\
& \times \exp \left( -\frac{1}{2} \left[ \sigma^2(T-t) - 2 \left( \log \frac{S_t}{y} + \left( r + \frac{1}{2}\sigma^2 \right) (T-t) \right) \right] \right) \\
= & \exp \left( -\frac{1}{2} \left[ -2 \log \frac{S_t}{y} - 2r(T-t) \right] - \frac{1}{2} \left( d_t^T(y) \right)^2 \right) \\
= & e^{r(T-t)} \frac{S_t}{y} \exp \left( -\frac{1}{2} \left( d_t^T(y) \right)^2 \right),
\end{aligned}$$

and

$$\exp \left( -\frac{1}{2} \left( -d_t^T(y) + 2r\sqrt{T-t}/\sigma \right)^2 \right)$$

$$\begin{aligned}
&= \exp\left(-\frac{1}{2}(d_t^T(y))^2 - \frac{1}{2}\left[\frac{4r^2}{\sigma^2}(T-t) - \frac{4r}{\sigma}d_t^T(y)\sqrt{T-t}\right]\right) \\
&= \exp\left(-\frac{1}{2}(d_t^T(y))^2\right) \\
&\quad \times \exp\left(-\frac{1}{2}\left[\frac{4r^2}{\sigma^2}(T-t) - \frac{4r}{\sigma^2}\left(\log\frac{S_t}{y} + (r + \frac{1}{2}\sigma^2)(T-t)\right)\right]\right) \\
&= \exp\left(-\frac{1}{2}(d_t^T(y))^2\right) \\
&\quad \times \exp\left(\frac{-2r^2}{\sigma^2}(T-t) + \frac{2r}{\sigma^2}\log\frac{S_t}{y} + \frac{2r^2}{\sigma^2}(T-t) + r(T-t)\right) \\
&= e^{r(T-t)}\left(\frac{S_t}{y}\right)^{\frac{2r}{\sigma^2}}\exp\left(-\frac{1}{2}(d_t^T(y))^2\right).
\end{aligned}$$

□

Similar calculations using (4.5.1) are possible for other Lookback options, such as options on extrema and partial Lookback options, cf. [Kha02].

*Acknowledgments.* We thank the referee for a careful reading of this paper, and E. Bichot for contributing to the simulation graphs. The results of this paper were presented in part at the AMS-Scandinavian meeting, Odense, 2000.

## References

- [AOPU00] K. Aase, B. Øksendal, N. Privault, and J. Ubøe. White noise generalizations of the Clark-Haussmann-Ocone theorem with application to mathematical finance. *Finance and Stochastics*, 4(4):465–496, 2000.
- [Bel99] N. Bellamy. Evaluation et couverture dans un marché dirigé par des processus discontinus. Thèse, Université d’Evry, 1999.
- [Ber98] H. Bermin. Essays on Lookback options: a Malliavin calculus approach. PhD thesis, Lund University, 1998.
- [Cla70] J.M.C. Clark. The representation of functionals of Brownian motion by stochastic integrals. *Annals of Mathematical Statistics*, 41:1281–1295, 1970.
- [DJ98] R.-A. Dana and M. Jeanblanc. *Marchés financiers en temps continu*. Economica, Paris, 1998.
- [DP99] M. Dritschel and Ph. Protter. Complete markets with discontinuous security price. *Finance and Stochastics*, 3(2):203–214, 1999.
- [É90] M. Émery. On the Azéma martingales. In *Séminaire de Probabilités XXIII*, volume 1372 of *Lecture Notes in Mathematics*, pages 66–87. Springer Verlag, 1990.

- [JP02] M. Jeanblanc and N. Privault. A complete market model with Poisson and Brownian components. In R. Dalang, M. Dozzi, and F. Russo, editors, *Seminar on Stochastic Analysis, Random Fields and Applications (Ascona, 1999)*, volume 52 of *Progress in Probability*, pages 189–204. Birkhäuser, Basel, 2002.
- [Kha02] Y. El Khatib. Contributions to the study of discontinuous markets via the Malliavin calculus. Thèse, Université de La Rochelle, 2002.
- [KO91] I. Karatzas and D.L. Ocone. A generalized Clark representation formula with application to optimal portfolios. *Stochastics and Stochastics Reports*, 34:187–230, 1991.
- [LL96] D. Lamberton and B. Lapeyre. *Introduction to stochastic calculus applied to finance*. Chapman & Hall, London, 1996.
- [Mey89] P.-A. Meyer. Construction de solutions d’ “équations de structure”. In *Séminaire de Probabilités, XXIII*, volume 1372 of *Lecture Notes in Math.*, pages 142–145. Springer, Berlin, 1989.
- [Øks96] B. Øksendal. An introduction to Malliavin calculus with applications to economics. Working paper no. 3, Institute of Finance and Management Science, Norwegian School of Economics and Business Administration, 1996.
- [Pha00] A. Phan. Contributions étude des équations de structure markoviennes. Thèse, Université de Strasbourg, 2000.
- [Pri96] N. Privault. On the independence of multiple stochastic integrals with respect to a class of martingales. *C. R. Acad. Sci. Paris Sér. I Math.*, 323:515–520, 1996.
- [Pro90] Ph. Protter. *Stochastic integration and differential equations. A new approach*. Springer-Verlag, Berlin, 1990.
- [Pro01] Ph. Protter. A partial introduction to financial asset pricing theory. *Stochastic Process. Appl.*, 91(2):169–203, 2001.
- [PSV00] N. Privault, J.L. Solé, and J. Vives. Chaotic Kabanov formula for the Azéma martingales. *Bernoulli*, 6(4):633–651, 2000.
- [Tav99] G. Taviot. Martingales et équations de structure : étude géométrique. Thèse, Université de Strasbourg, 1999.
- [Yor76] M. Yor. Sur les intégrales stochastiques optionnelles et une suite remarquable de formules exponentielles. In *Séminaire de Probabilités, X (Seconde partie: Théorie des intégrales stochastiques, Univ. Strasbourg, Strasbourg, année universitaire 1974/1975)*, pages 481–500. Lecture Notes in Math., Vol. 511. Springer, Berlin, 1976.